


Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Cusanty european Caule filiformi, glomendis capitatio sessibles

floribus singulis brevissime pediallates 4-5 partitos

tubo corollae applit subrito relytis segmente ovata oblusitisonle et. et lacrimies limbi obbusiusenlas erectus demen fatentes, aequante; stamphibus linke brevioribus unthuis ovalis squamis missionis basi vosoble presentes, aprice tentations findritte, adpressis, stylis sumastignately distanium gerbowedynnessen aegerantibes divergentibus, stymatibis skylis mulle long britis gapsula globosa depressa (s. seboonia) britis corolla marres cente calyptanta. Europe Or Coylus Ultisz , bannelig thumles sto. _ Shop Thown has oblected a variety near Continue with a rather antish ovary but apparently not difference in structure for the common form, as the Mannen proies with a stylopodi- do) more erect styles, and a somewhite comir saprule. Lean find no other difference in the Therian Cylono me The plant is larger as the forces C.En. En the stems stouter (certainly not regullar) as Chorsey fund sags ber pag. 173) the flowers of about the same size but alex one routh obtained, and small, freeze small largen al rowlle fire tom from it bese as covering only the top of the capsule. The own whith in the other speries only takes a very some. spacewithing the tide of the cowlly fill it very soon aftermeds to that the corolle appears unevelle, and the same Time much longer than the alex which is only the consequence of its line either afrom its inserted on top of the growing capsale.




Cusaula Caule rapillares vamoris imo glomendes floribusque singulis arate sessilibus floribus 4 partitos tubo corollae planeviter cumpanulato algois appulati segmentis transvers ovatis Carpidatis s. abrupte aruminatis plessy lavinis flessy lavinis limbi aruminatis breviore Stamme bus limbo Dien Sio brevioribus. squaring late ovatis crenatis s. brevite fundrialis, sometiments or convergent bus?), shylis oraniem (atropurous) stylorum long. Tucke, Mirabello
on some shrubby Labiata, a smill Thymas? Howers smallen Thus on any other species Thour to me; Istinguish a besile, by it short openly campa unlate tube of the corolla, flower, - the spensku exammed by me teller mone the A partie but frequently only 3 parties only x large in proportion to the corolla and not adpressed more compulate lobes carrinate with The lobes of the corolla are top arite tiberrely or acuminate the influenced at the finh a o just below it somewhat sacrate so that they appear as with a little Elut sac or spun on the under or outer side. - Stymas in my spenten dack purple and as long as the tyle forpen but thicker, and blunde appearing to be The tyle a clavate appearance to I observed 3 styles. The sowllawille will former of the apple The name minima would be propen Near C. Chithy man from Shirts it Siffers & the Vi partir flowers by the by the broad lobes of the calyx a some by the short openly campandet the of still more closely allied it to Combingtony smill parter flowers for which is is is the spreading the stand the smile for the spreading the corolla the corolla as by the short styles; apparelly

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN



Teres Cusinti ambien Caule orgillerer, He glomestis capitation Tenilibus, som len mynlis brevis. me pedirelletis, 4-5 partitis, tube coulle campunelete algors segmentis overtig -. ovite-landolatis acentis breviore Cavinias limbs ovitas acentes applicatel inflexas margine levissime oremeletes, patantes aequante, slaminibres subulation limbo brevioribres, an theris cordato- Triangulanty squames quadritio apire trumato findrites, temissimo, tubo molhe brevioribus, Aplis brevisimis omne sligmitibus ovarium globosum subrequestion paule Dongentitus; magasala corolla (177mm ompsulam globorane stylis horizetalit Diregentitus conventaces at the prostate marcescente.

De Arabig (Shimpun, Mystac)

1835. Agypto (vide Chining monogn.) et Aleysinin (Withingung 1843, on Labitis, G. brevisty la A. am) This species very nearly resembles C. Epithy mum, but it is well distriquished by the way short obyles being much shorter than the stry mas as with together gust equally the orang the ly the finely cremeted manying their square soules. by the finely cremeted manying their square soules. by the sather triangular authors and more subulate stilaments. by The larger. ovary which fills the whole the of the moble, fine this respect, AR in the sorles, also in The quadriputes onle being fregantly to parter it resembles me Coura, cumprien, but the shole flower ar expensely the rappule is much smillen, and emplule a sudd are of the same sign as Co Epithyun The Ambian sperimen before me has Carger flowers with the the Californial not much exceeding the state of the world

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN


MISSOURI
BOTANICAL
GARDEN

x as not so had present, the obystiman specime is, in full it has therefore quite adfent materity of fruit, corollar agrants, appearance. I smallen not much exceed the color its lobes more reflexed an something result to the position and capsale covered to the base to course of and the papearance of the base of the separates from a confidence of the base of the ba MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS MISSOURI BOTANICAL copyright reserved

GARDEN

cm

Cusanta Epilinum Weihe Caule Simplier capi blacco), glomendis ante sessitibus, floribus singulis sessitibus se partitus tralge is campanaletti, impulati, confluent glibrand segmentis ovates autim prosiculation acutions. abrupte avuminatis, vorolla globoza relgeen vix excedente, tubo lavinias ovatas acutas incurvos subregiante, staminibis limbos Simbos Similio Freviori bus filamenti; brevisionis; squaris affatusouratis tournatos trevita findriat; plennyne) Saucem corolla attitionation adpression plening stylis run stigmation of vorario globoso (my mo) immistis stylis brevissius deple longionitus corolla orra On Linum usitetissimum in German Pennyhomi - (Sange) France, lagland, also in Massa husats (Jakes) This species is distinguished for most others by its simple steen and by its strictly ashering to and species of plants. Dam at less not aware of its having even been for a any where else but on flax. The flowers the very compact globone glowenly the caly , snortly sparted but sell elso 6 and 4 parties, - The first water I fart it with 3 styles . _ it loke, are invenced so that not spread the stance included the world the better the followed but The could in the of the world is the followed but The could in the more or con make with the scales in my garmen sperimens want of Gronois. The scales in my game sperimens are fly as large as the tall a I am slightly fundate; in english and they are shorter auen men spenden and irregularly toother at the townste Lorens apex, or occasion bilobed. Stymas thirtee that wery The over is large, - proportion as Jils the whole of the worlde After flowers the the corolle is not tom form its insertion, but the property



Canationsis

Caule capillace tencissino. parmoso.

glomendis arote sersilibus, parcifloris, floribus,

singulis sersilibus, 5 partitis; requis

capulati regrentis volus acentisus calis

tubs corollae companulato segmentis calgoris

cipulati contis acentis ralis braviore, laciaias

cindi ovatas acentas erectas de mun petentes

cindi ovatas acentas erectas de mun petentes

bravioribus; squanito latis, tencissimis

api ce vorene leto-findriatis, stylis cum

stigment bus Sivariatis ovarium globosum

aequantibus, corolla circum appulace

globoso-depressam massesset as basis

jersistente

Tenerifa _ Hart Arav Philas sub nom C. EpiG'num

Very near C. Ejer Cinum but flowers and glomentes only half is large stems repicling a se branching which is never the cree - C. Ejer.

and, cally not involuding the corolle as much of corolle of corolle of much a faith of the compared, I have seen it only with marky ripe print much only with marky ripe print much only with

MISSOURI BOTANICAL GARDENT GEORGE ENGELMANN PAPERS



the corollee ampanulato subsequento calgois segments ovals acien kate pul pleningue excedente lacinias litulio ovatas accumulas de crectas si patentes demum reflexas sulla limbo sulla resolución de acquento staministrationes sulla limbo brevión bey anthem ovata orolla borcosonto ovatas cremulatios tubo conolla borcosonto ovarian globosium deplo superatitions apir solumnodo di stymatibus stylos aequentitues pale vergentitues pele vergentitues pele vergentitues pele vergentitues from the man resent file dente a bisi plening diversity France, Italy & the Pyrenes In mendous, highly of other with Ay In The perplash A small plant, with capillacever stens and compact clusters of acute flowers these acute lobis destruguesh it at once from County actions as also the Shorter file to the long its of the sound inore erect is very their members - other somewhat the ker and john clongation gland or tuberoles it is who test or red of he are the stymas es authus are also sometime Deck parple, some true pele alex paler or yellowsh. For flowers ene when you they other wines the Colors of from my spiritures the aly & Corolla were Littyry Lotter, Implie. Med rago, genista, partien of the light of the series of the s also on composité, crepis, Alles, Centeures Sematula althouse the town from its buse Arbij Clea, Chrysanthem ,- on Labratec. and somewhat elevate ... Lavalle Forther Also on Callenia The mande of not helpas large Erica Tommelelle Liamon galier was found they such and on Grasses (School) dasable layer to other places and on Grasses (ESS) and - Equirate (De Cham) Ship as an factor of the such as a factor of th frontleren event amongst The Erica of the Cape of good hope in the Carlon

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

MISSOURI

BOTANICAL

GARDEN

bearded grape, The barbata, figured strend by Tabernaemontanes (p. 1287) haf ite and their period this Currents of the grape vine whene it is of steinle as hangs down from the grupe i long opiller (sometias a jost la capillery filaments, somewhat recember) a bean. B. pans flom A vaniety with much smaller flervers and aruminate lobes of the calgy which are in longer than e the the toute on the loves of the corollar, grows near Friest floorger on Satureja Inontan. (Benjer in Herb A. Braun) The same variety only Afternay of the light wider lober of the Baky x or runs on Lavadla Slovehar i Th Syrences (: Ht. A. Braun). In Soft the appele is very small and versule I se envelloper by the corolla which the des not exactly over the a supplied fort. It some for does at sometimes over with a large city which well to the total is as try of the take of the world MISSOURI BOTANICAL GARDENS
GEORGE ENGELMANN PAPERS MISSOURI

BOTANICAL copyright reserved GARDEN


Cluten Cascula Caule capillarco vamossissimo glomembis bratistis, and sessible segmentis lancelle acuminatis demellas punto brevionila the corolle brevitis en imparentato segment, carrie segment, carrier segment, orders segments Dimites Grevione: staminibus butante Cinho J. Zu. Si Crenty; mædim skingestilm, authur, ovati. squares bifiles ? faceren the, the addingention; sky e; ovaris (globon ?) vjekt longe solle repole : Forther Tyrol on Colutes artores Tyrol castle . H. A. Braun' num Fynderatte My sperimen is just beginning to flower flower, but it appears to be well to character of the large only x with a sungland in the last of x with a sungland in the last of the corolla, the laber of their are othere said the only flower som by he event but por bibly spreading afternal, the senter are as long as the short tube, but appear not to be fringer or overate but if sam not much mis taken bifile! At the base of the closely sensely joing glomentes large water warming brack, and compriences, The flowers vesemble somewhit Those of C. Cuithy men by the accuminate lobes of the call X: en are welly diffinguished of the shape of the small corolla l Les approuls somewhy will protably be found at base

BOTANICAL

GARDEN

copyright reserved

cm



Cascula cupulata ". 1. Cente capillarco, glomerulis multiflons capitati ante sessibles, floribes singulis ante sessibibles sub-s partitis, calyons campanulate enpulate lobis brevisionis

transvers: - ovatis, al supte cuspodatis mis testen Itim algundong to Since the corollae comparaulito orly ce brevione

segment ord lavinias orates actificandas erectas Somen patentes aequante stamshiles Combo brevion been anther of the osbirelety squamis truncation aprice findication to the selle velex det tes i convergentites. attingent, ty stylis rum stigmettetes ovarrium seprem_ globes um Bupements bus, aprice Divingentilus, Aigmetilus olyles aguantibus, somble orsner apsalam globosam cirrumscissam munescente non elemente paulo a besi nom Alter, my sperimens of Artemisis and medings? coll by Lebour. This species is way near to Epithy mun from this species by its early & shih covers the greater which is mearly entire and it that part of the flower respect similar to C. Balylomia finth from forth' only is de it is devided it. Allei) & the bond abtuse bles which are very shortly amunite or cresporte The loke of The whole flower is els. The C. Gutty me smallen the aprole but half so leve the lobes of the corolla not accuminate the singly would generally allow the level of the ord, lavinite at the townstaper. ripe capsule, at is not as in (Epithyn and appears to be attributed to it at is base the place when il reparates for its bree, in C. Epitym is is raised higher up by the entargents. of the capsule and approaches more (European where it only lovers the top of Capsule hoodlike MISSOURI


BOTANICAL copyright reserved cm GARDEN or divided nearly to the middle (in the specimens)

for the Taurian deart), the lobes would internated and
in all other respects similar to the other form MISSOURI BOTANICAL GARDEN.

GEORGE ENGELMANN PAPERS MISSOURI BOTANICAL

copyright reserved cm GARDEN

Cusanta oupalata caule sufficient, glomerules multiflons capitati semilibus. floribus Tingulisant sessilibies 5 partitis, only vis campanulity capulati lobis obovato orbivalatis glandulose carinatis quia subvaspidatis teles conolle Parinias limbi ovales abbusas enetas Jemum patente, æguante, stanin, En timbe brevioriber antheris orbitality symmes ovatis event fishates toben vorder attrujentibus vonvergentitus Appliseum skjænteten ovanium denne globore din supermile (Jonatoly Styles) subarquantitus. crops els Or the Terek near Kristan i, Cirmeria This popula that this is only a variety of Comments. Isling wisher by touten stein and branched strongen texture of the cely x and cowler but expensely bolluse lobes of coulle and shorter styles. I have not seen ang flowers more et vænned um ang søfe fruit.



H.B. F. 1825 GEORGE ENGELMANN PAPERS MISSOURI BOTANICAL CARDEN MISSOURI BOTANICAL copyright reserved

GARDEN

cm

Casouta Besquamate ny. Care filigomi glomendis sessilibus Cax is floribus singulis pedrellatis sul 4 partitis, tube corollee cumpa mulato calgoris sogmentes ovato tisang Carilies sularutis brevivre, lavinia l'imbi ovato Camedatas deternas erectas aequante, staminibus limbs Suis. brevionibus, antheris ovates, squamis carentibus, stylis ovanium globorum, Conjonibus Siani vix aequantibus. " orolle ad a in arrum Apples Sivarientes covornatain adein manufacture HG. Aug Thies sub nomine C. Epithymum, on Ranummler ? - Oblgsom Duneton? This species belongs to the section of Contre for the old world will a lange olyx a smill this mention Corolla (C. Epilinum C. planiflore ambica.
C. minuta & Tuppellate) esperially near The two first ones; and Satingent for all by the stouter not all by absence of the bulfila menton, and principally soules! Amongot many flowers example I forme only in on a trave of a scale i the form of two leeth at the laxe of one of the filaments. The flowers are 4 and very morely 5 parter, - the large rather firm orly includes towers the greater part of the corolla the fiturnat being only half as land as the exect and even a little inflexed lokes dinte of the conolle are little seen. the globose overy is rather large in proportion and fills the whole with the of the milla. It size - 2 to short Tyles Sisting with the Car this speries to which it appear to bear the close affinity.